



Mitochondria in Aging and Disease: From Cellular Powerhouse to Pathological Trigger

Vandana Patil*¹ Mangala Thakare²

*¹Department of Zoology, Kamla Nehru Mahavidyalaya, Sakardara, Nagpur, Maharashtra, India

Patil2vandana@gmail.com¹

²Department of Zoology, G. K. Art and Science Mahila Mahavidyalaya, Kawarabandh, Tah. Salekasa, Gondia,

mangu81084@gmail.com²

Abstract

Mitochondria are multifunctional organelles that play an essential role in cellular energy production and metabolic regulation. Traditionally recognized as the “powerhouses” of the cell, mitochondria are now known to participate in numerous biological processes including apoptosis, calcium homeostasis, reactive oxygen species (ROS) generation, and regulation of cellular signalling pathways. Increasing evidence suggests that mitochondrial dysfunction contributes significantly to the aging process and the development of several age-related diseases such as neurodegenerative disorders, cardiovascular diseases, metabolic syndromes, and cancer. Aging is a complex biological process characterized by the progressive deterioration of physiological systems, resulting in impaired function and a higher risk of death. It is associated with structural and functional alterations in mitochondria, including mitochondrial DNA mutations, impaired oxidative phosphorylation, increased ROS production, and decreased mitochondrial quality control mechanisms. These alterations disrupt cellular homeostasis and accelerate tissue degeneration. This review summarizes current knowledge on mitochondrial structure and function, the mechanisms responsible for mitochondrial dysfunction during aging, and their involvement in age-related diseases. Additionally, potential therapeutic strategies aimed at preserving mitochondrial function are also discussed.

Keywords: Mitochondria, Aging, Mitochondrial dysfunction, Oxidative stress, Age-related Diseases

1. Introduction

Aging is a complex biological process characterized by progressive decline in cellular and physiological functions, which ultimately increases susceptibility to chronic diseases. Several molecular mechanisms have been proposed to explain the aging process, including genomic instability, telomere attrition, epigenetic alterations, and mitochondrial dysfunction (López-



Otín *et al.*, 2013). Among these factors, mitochondrial dysfunction has emerged as one of the most significant contributors to cellular aging and age-associated disorders.

Mitochondria are double-membrane organelles responsible for producing cellular energy in the form of adenosine triphosphate (ATP) through oxidative phosphorylation (Wallace, 2005). In addition to energy production, mitochondria regulate multiple cellular processes such as apoptosis, redox signalling, calcium homeostasis, and metabolic regulation (Green & Reed, 1998; Lane, 2011). Because of their central role in maintaining cellular homeostasis, mitochondria are often considered key determinants of cell survival and function.

Unlike other cellular organelles, mitochondria possess their own genetic material known as mitochondrial DNA (mtDNA). This genome encodes essential components of the electron transport chain required for ATP production. However, mtDNA is highly susceptible to oxidative damage due to its proximity to the electron transport chain and the absence of protective histone proteins (Bratic and Larsson, 2013). As a result, mutations accumulate in mitochondrial DNA over time, which can impair mitochondrial function and contribute to cellular aging.

Another important feature of mitochondria is their dynamic nature. Mitochondria constantly undergo fusion and fission processes that allow them to maintain their structural integrity and adapt to metabolic demands (Sun *et al.*, 2016). These processes are essential for maintaining mitochondrial quality control. However, disruptions in mitochondrial dynamics can lead to the accumulation of damaged mitochondria within cells.

Accumulating evidence indicates that mitochondrial dysfunction is closely associated with the aging process. Age-related mitochondrial changes include decreased ATP production, increased generation of reactive oxygen species, reduced mitochondrial biogenesis, and impaired mitophagy (Balaban *et al.*, 2005; Sun *et al.*, 2016). These alterations compromise cellular metabolism and promote the development of degenerative diseases.

Mitochondrial dysfunction has been implicated in numerous age-related diseases including Alzheimer's disease, Parkinson's disease, cardiovascular diseases, metabolic disorders, and cancer (Wallace, 2005; López-Otín *et al.*, 2013). Because mitochondria are involved in such a wide range of cellular processes, maintaining mitochondrial health is essential for preserving cellular function during aging.

Therefore, understanding the mechanisms underlying mitochondrial dysfunction may provide valuable insights into the aging process and help identify potential therapeutic targets for preventing age-related diseases.



2. Structure and Function of Mitochondria

Mitochondria are highly specialized organelles characterized by a double-membrane structure consisting of an outer mitochondrial membrane and an inner mitochondrial membrane. The outer membrane functions as a protective barrier and contains protein channels that regulate the transport of metabolites and ions between the mitochondrion and the cytoplasm (Lane, 2011).

The inner mitochondrial membrane is highly folded into structures called cristae, which increase the surface area available for biochemical reactions involved in oxidative phosphorylation. Embedded within this membrane are the protein complexes of the electron transport chain that generate ATP through a series of redox reactions (Wallace, 2005).

The mitochondrial matrix contains enzymes involved in the tricarboxylic acid (TCA) cycle, mitochondrial DNA, ribosomes, and various metabolic intermediates necessary for cellular respiration (Bratic & Larsson, 2013).

Mitochondria perform several essential functions in the cell, including:

- Production of ATP through oxidative phosphorylation (Wallace, 2005)
- Regulation of programmed cell death or apoptosis (Green & Reed, 1998)
- Maintenance of calcium homeostasis (Balaban *et al.*, 2005)
- Generation and detoxification of reactive oxygen species (Balaban *et al.*, 2005)
- Regulation of metabolic pathways including fatty acid oxidation and amino acid metabolism (Lane, 2011)

Due to these diverse functions, mitochondria are considered central regulators of cellular metabolism and survival.

3. Mitochondria and the Biology of Aging

The relationship between mitochondria and aging has been widely studied in recent decades. One of the most influential theories explaining this relationship is the mitochondrial free radical theory of aging, which suggests that the accumulation of oxidative damage caused by reactive oxygen species contributes significantly to the aging process (Balaban *et al.*, 2005).



During normal mitochondrial respiration, electrons are transferred through the electron transport chain to generate ATP. However, a small fraction of these electrons may leak and react with oxygen to form reactive oxygen species such as superoxide radicals. Although cells possess antioxidant defence mechanisms to neutralize these molecules, excessive ROS production can cause damage to cellular components (Sun *et al.*, 2016).

Over time, oxidative damage accumulates in mitochondrial proteins, lipids, and DNA, leading to impaired mitochondrial function and reduced energy production (Bratic & Larsson, 2013). Furthermore, damaged mitochondria may generate even higher levels of ROS, creating a cycle of oxidative stress that accelerates cellular aging.

Another important factor contributing to mitochondrial aging is the decline in mitochondrial biogenesis. Reduced activity of transcriptional regulators such as PGC-1 α can limit the formation of new mitochondria, thereby decreasing cellular energy capacity (López-Otín *et al.*, 2013).

4. Mitochondrial Dysfunction in Age-Related Diseases

Mitochondrial dysfunction is strongly associated with several chronic diseases that commonly occur during aging. Neurodegenerative diseases such as Alzheimer's disease and Parkinson's disease are characterized by impaired mitochondrial energy production and increased oxidative stress, which ultimately lead to neuronal degeneration (López-Otín *et al.*, 2013). Similarly, mitochondrial dysfunction in cardiac cells can impair energy metabolism and contribute to cardiovascular diseases including heart failure and ischemic injury (Wallace, 2005). Metabolic disorders such as obesity and type 2 diabetes are also associated with mitochondrial dysfunction. Reduced mitochondrial oxidation of fatty acids and glucose contributes to metabolic imbalance and insulin resistance (Bratic & Larsson, 2013).

In cancer, mitochondrial alterations influence metabolic pathways that allow tumor cells to survive and proliferate rapidly under altered metabolic conditions (Wallace, 2005).

5. Mitochondrial Quality Control Mechanisms

To maintain cellular homeostasis, mitochondria possess several quality control mechanisms that regulate their structure, number, and function. These mechanisms include mitochondrial biogenesis, mitochondrial dynamics (fusion and fission), and mitophagy. Proper regulation of these processes ensures that damaged or dysfunctional mitochondria are removed and



replaced with newly synthesized organelles. Mitochondrial biogenesis is regulated by transcriptional coactivators such as peroxisome proliferator-activated receptor gamma coactivator-1 alpha (PGC-1 α), which stimulates the expression of nuclear and mitochondrial genes required for mitochondrial replication and function (Scarpulla, 2011). Aging is often associated with decreased expression of PGC-1 α , leading to reduced mitochondrial biogenesis and impaired cellular energy metabolism.

Mitochondrial dynamics involve continuous cycles of fusion and fission that maintain mitochondrial integrity. Fusion allows mitochondria to mix their contents and dilute damaged components, while fission facilitates the removal of damaged mitochondria through mitophagy (Youle & van der Bliek, 2012). Disruption of these processes can result in mitochondrial fragmentation and dysfunction. Mitophagy is a specialized form of autophagy that selectively removes damaged mitochondria. The PINK1-Parkin pathway plays a crucial role in this process by identifying and targeting defective mitochondria for degradation (Pickrell & Youle, 2015). Age-related decline in mitophagy leads to the accumulation of dysfunctional mitochondria and contributes to cellular aging.

6. Role of Mitochondrial Dysfunction in Specific Age-Related Diseases

6.1 Neurodegenerative diseases

Mitochondrial dysfunction has been strongly implicated in neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease, and Huntington's disease. Neurons are highly dependent on mitochondrial ATP production for maintaining synaptic transmission and neuronal signaling. Impaired mitochondrial function can lead to reduced energy supply, increased oxidative stress, and neuronal degeneration (Lin & Beal, 2006).

In Alzheimer's disease, mitochondrial abnormalities include decreased cytochrome oxidase activity, increased oxidative damage, and altered mitochondrial dynamics (Swerdlow, 2018). Similarly, mutations in genes involved in mitochondrial quality control, such as PINK1 and Parkin, have been linked to Parkinson's disease (Pickrell & Youle, 2015).

6.2 Cardiovascular Diseases

The heart requires a continuous supply of ATP to maintain its contractile activity. Approximately 90% of cardiac ATP is generated by mitochondria through oxidative phosphorylation. Mitochondrial dysfunction can therefore severely impair cardiac energy



metabolism and contribute to conditions such as cardiomyopathy, ischemic injury, and heart failure (Rosca & Hoppel, 2010).

Excessive ROS production in cardiac mitochondria can damage mitochondrial membranes and proteins, leading to reduced ATP synthesis and impaired cardiac function.

6.3 Metabolic Disorders

Mitochondrial dysfunction also plays a critical role in metabolic diseases such as obesity and type 2 diabetes. Reduced mitochondrial oxidative capacity in skeletal muscle and adipose tissue can impair glucose metabolism and promote insulin resistance (Lowell & Shulman, 2005).

In addition, altered mitochondrial fatty acid oxidation may lead to lipid accumulation in tissues, which further contributes to metabolic dysfunction.

6.4 Cancer

Cancer cells exhibit significant alterations in mitochondrial metabolism. Although many cancer cells rely heavily on glycolysis for energy production, mitochondria remain essential for biosynthetic processes and metabolic signaling (Vyas *et al.*, 2016). Mutations in mitochondrial DNA and alterations in mitochondrial metabolic pathways can promote tumor growth and survival.

7. Therapeutic Strategies Targeting Mitochondrial Function

Recent advances in biomedical research have led to the development of therapeutic strategies aimed at improving mitochondrial health and preventing age-related diseases. Several mitochondria-targeted antioxidants, such as MitoQ and SkQ1, have been developed to reduce oxidative stress within mitochondria (Smith & Murphy, 2010). These compounds selectively accumulate within mitochondria and neutralize reactive oxygen species. Another promising approach involves stimulating mitochondrial biogenesis through activation of PGC-1 α signaling pathways. Lifestyle interventions such as regular physical exercise and caloric restriction have also been shown to enhance mitochondrial function and improve metabolic health (Hood *et al.*, 2019).



Furthermore, pharmacological agents that stimulate mitophagy may help remove dysfunctional mitochondria and restore cellular homeostasis.

8. Future Perspectives

Although significant progress has been made in understanding mitochondrial biology, many aspects of mitochondrial involvement in aging remain unclear. Future research should focus on identifying molecular pathways that regulate mitochondrial quality control and determining how these pathways change during aging. Advances in genomic and proteomic technologies are expected to provide new insights into mitochondrial signaling networks and their role in disease development. Additionally, the development of mitochondria-targeted therapies may open new avenues for treating age-related disorders and promoting healthy aging.

Table 1. Major Age-Associated Diseases Linked to Mitochondrial Dysfunction

Disease	Mitochondrial Mechanism	Key Effects	References
Alzheimer’s disease	mtDNA mutations, impaired OXPHOS	neuronal degeneration	Swerdlow, 2018; López-Otín et al., 2013
Parkinson’s disease	defective mitophagy (PINK1/Parkin)	dopaminergic neuron loss	Pickrell & Youle, 2015
Cardiovascular diseases	ROS production	endothelial dysfunction	Wallace, 2005; Rosca & Hoppel, 2010
Diabetes mellitus	impaired mitochondrial metabolism	insulin resistance	Lowell & Shulman, 2005; Bratic & Larsson, 2013
Cancer	metabolic reprogramming	uncontrolled proliferation	Vyas et al., 2016; Wallace, 2005

Table 2. Key Mitochondrial Quality Control Mechanisms



Mechanism	Key Proteins	Function	References
Mitochondrial fusion	Mfn1, Mfn2, OPA1	maintain mitochondrial integrity	Youle & van der Bliek, 2012; Sun et al., 2016
Mitochondrial fission	Drp1	remove damaged mitochondria	Youle & van der Bliek, 2012; Sun et al., 2016
Mitophagy	PINK1, Parkin	degradation of dysfunctional mitochondria	Pickrell & Youle, 2015
Biogenesis	PGC-1 α , NRF1, TFAM	formation of new mitochondria	Scarpulla, 2011; López-Otín et al., 2013

Table 3. Potential Therapeutic Strategies Targeting Mitochondria

Strategy	Example	Mechanism	Reference
Antioxidants	CoQ10, MitoQ	reduce oxidative stress	Smith & Murphy, 2010
Caloric restriction	dietary intervention	enhances mitochondrial function	Hood <i>et al.</i> , 2019
Exercise	endurance training	stimulates mitochondrial biogenesis	Hood <i>et al.</i> , 2019
NAD ⁺ boosters	NMN, NR	improve cellular metabolism	Balaban <i>et al.</i> , 2005; Sun <i>et al.</i> , 2016

9. Conclusion

Mitochondria are central regulators of cellular metabolism, energy production, and signaling pathways. During aging, mitochondrial function gradually declines due to oxidative stress, mitochondrial DNA mutations, impaired mitochondrial dynamics, and reduced mitophagy. These alterations contribute to the development of numerous age-related diseases including



neurodegenerative disorders, cardiovascular diseases, metabolic syndromes, and cancer. Understanding the molecular mechanisms underlying mitochondrial dysfunction may provide valuable insights into the aging process and facilitate the development of therapeutic strategies aimed at preserving mitochondrial health and improving human longevity.

References

1. Balaban RS, Nemoto S, Finkel T. 2005. Mitochondria, oxidants, and aging. *Cell*. 120:483-495.
2. Bratic I, Larsson NG. 2013. The role of mitochondria in aging. *Journal of Clinical Investigation*. 123:951-957.
3. Green DR, Reed JC. 1998. Mitochondria and apoptosis. *Science*. 281:1309-1312.
4. Lane N. 2011. Mitonuclear match: optimizing fitness and fertility over generations drives aging within generations. *BioEssays*. 33:860-869.
5. López-Otín C, Blasco MA, Partridge L, Serrano M, Kroemer G. 2013. The hallmarks of aging. *Cell*. 153:1194-1217.
6. Sun N, Youle RJ, Finkel T. 2016. The mitochondrial basis of aging. *Molecular Cell*. 61:654-666.
7. Wallace DC. 2005. A mitochondrial paradigm of metabolic and degenerative diseases, aging, and cancer. *Science*. 283:1482-1488.
8. Hood DA, Memme JM, Oliveira AN, Triolo M. 2019. Maintenance of skeletal muscle mitochondria in health, exercise, and aging. *Annual Review of Physiology*. 10:81:19-41.
9. Lin MT, Beal MF. 2006. Mitochondrial dysfunction and oxidative stress in neurodegenerative diseases. *Nature*. 19;443(7113):787-795.
10. Lowell BB, Shulman GI. 2005. Mitochondrial dysfunction and type 2 diabetes. *Science*. 21;307(5708):384-387.
11. Pickrell AM, Youle RJ. 2015. The roles of PINK1, Parkin, and mitochondrial fidelity in Parkinson's disease. *Neuron*. 21;85(2):257-273.
12. Rosca MG, Hoppel CL. 2010. Mitochondria in heart failure. *Cardiovascular Research*. 88(1), 40-50.
13. Scarpulla RC. 2011. Metabolic control of mitochondrial biogenesis through the PGC-1 family. *Biochimica et Biophysica Acta*. 1813(7):1269-1278.
14. Smith RAJ, Murphy MP. 2010. Mitochondria-targeted antioxidants as therapies. *Trends in Pharmacological Sciences*. (57):106-114.



15. Swerdlow RH. 2018. Mitochondria and mitochondrial cascades in Alzheimer's disease. *Journal of Alzheimer's Disease*. 62(3):1403-1416.
16. Vyas S, Zaganjor E, Haigis MC. 2016. Mitochondria and cancer. *Cell*. 166(3);555-566.
17. Youle RJ, van der Bliek AM. 2012. Mitochondrial fission, fusion, and stress. *Science*. 31;337(6098):1062-1065.